



# ORCHARD FACTS



February 23, 2012  
Vol. XIII, No. 1

## In This Issue

Walnut Blight Management

Mobile Irrigation Lab

Training and Pruning Young  
Walnut Trees

Tools for Weed Management  
Decisions

Walnut Pruning Field Meeting

Bill Krueger  
Farm Advisor

In accordance with applicable State and Federal laws and University policy, the University of California does not discriminate in any of its policies, procedures, or practices on the basis of race, religion, color, national origin, sex, marital status, sexual orientation, veteran status, age, medical condition, or handicap. Inquiries regarding this policy may be addressed to the Affirmative Action Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6th Floor, Oakland, CA 94612-3560. (510) 987-0097.



To simplify information, trade names of products have been used. No endorsement of named products is intended, nor is criticism implied of similar products which are not mentioned.

*Sacramento Valley Walnut Newsletter*

## Walnut Blight Management

*Richard P. Buchner – UC Farm Advisor, Tehama County*

*Steve E. Lindow – Professor of Plant Pathology, UC Berkeley*

Our most current information indicates the section 18 for ethylene bis-dithiocarbamate materials (Manzate flowable or Manzate Pro-stick) for walnut blight control are on track with issuance expected about March first. We have not seen the actual 2012 section 18 but expect similar wording and restriction as in previous years.

Copper tank mixed with Manzate flowable or Pro-stick is currently the most effective spray choice. Good quality copper products are all effective for controlling walnut blight. Follow label rates because metallic rates and copper availability vary depending upon product. Full coverage at full material rates is recommended.


Walnut Blight (*Xanthomonas arboricola pv juglandis*) bacteria over winter in the outer bud scales or cataphylls. Within the dormant bud, the inner leaf tissue and flowers are pathogen free. As the shoot grows through the infected outer bud scales, bacteria have the opportunity to move and infect developing leaves, shoots and flowers. Infection occurs when rainfall and/or wet conditions transport blight bacteria to developing tissue. The probability of infection depends upon how much pathogen exists on individual buds and environmental conditions favoring bacterial spread and infection. First walnut blight sprays are timed to coincide with early shoot emergence. This places a protective layer of bactericide on leaf tissue. If bacteria are splashed from the outer bud scales to developing shoots and flowers, the bactericide barrier prevents infection and subsequent blight lesions. Since all walnut shoots do not emerge at the same time, the first protective spray is applied when 40% of the shoots are elongating and before leaves expand. This is usually referred to as the “prayer” stage since the unfolded leaves resemble hands held in prayer. A second spray is applied about 7 to 10 days later and is designed to protect the remaining opening buds. Additional spray decisions are based upon measurements of infected buds, disease history, weather conditions and variety.

Many walnut growers wonder if they need to blight spray Chandler or other late leafing varieties. We have measured over 50% crop damage on Chandler walnuts when overwintering bud populations were high and spring weather favored disease. Conversely, we have measured little to no blight on Chandler walnut with low to zero bud population levels even when wet spring weather favored disease. Late leafing walnut varieties have less opportunity time to build high walnut blight populations in dormant buds. As a result, initial inoculum levels are low. This does not eliminate, but reduces the probability of disease incidence. A good late leafing strategy would be to apply the first two applications with the intention of maintaining low inoculum levels.

Bud pathogen information, disease history and weather conditions can be used to improve spray decision accuracy. The California Seed and Plant Lab, 7877 Pleasant Grove Road, Elverta, California, 95626 (916-665-1581) is available to evaluate dormant walnut buds for blight bacteria. The sampling technique is available at [cete.hama.ucdavis.edu](http://cete.hama.ucdavis.edu). Click on orchard crop, click on walnuts and scroll down to “sampling dormant walnut buds.”

## Disease Cycle for Walnut Blight caused by *Xanthomonas arboricola* pv *juglandis*

Richard P. Buchner and Steve E. Lindow



## Mobile Irrigation Lab

The Mobile Irrigation Lab (MIL) is a service of the Tehama County RCD that performs on-site evaluations of agricultural irrigation systems. The two key pieces of information the MIL provides to the grower are, the rate of water being applied to a crop, and the overall efficiency of the irrigation system. The goal of the MIL is to give growers an overall snapshot of their irrigation system, provide scheduling information, and give recommendations to help improve efficiency.

The MIL provides several pieces of information in a full 10 to 20 page report that includes; application rates (how much water is being applied), the system's efficiency or Distribution Uniformity (DU), suggested run times, soils descriptions with maps, pump efficiency data, and a written summary full of tips and suggestions. Based on these evaluations recommendations are made for system repairs and developing maintenance plans that can help improve system performance. Mobile Lab Technicians can then work with growers to develop irrigation management practices that are the most efficient and cost effective.

Through this service, growers gain tools that allow efficient system operation, which in turn can lead to reduced energy costs and water use while simultaneously increasing yields. Thanks to support from concerned agencies, MIL evaluations are performed at no cost to growers. To schedule an evaluation, contact Kevin Greer at 527-3013x102 or email at [kevin@tehamacountyrcd.org](mailto:kevin@tehamacountyrcd.org) .

## Training and Pruning Young Walnut Trees

*Carolyn DeBuse, Farm Advisor, Yolo & Solano Counties;*

*Janine Hasey, Farm Advisor Sutter & Yuba Counties;*

*and Bruce Lampinen, Extension Specialist, UC Davis*

Training and pruning young walnut trees is done in the first 5-6 years of the life of the tree. Traditionally it has been done in a modified central leader pruning style. The height of the first branch, the amount of branch thinning, number of scaffolds chosen and the amount of wood taken off when heading cuts are made differs for hedgerow (132+ trees per acre) and traditionally spaced orchards (e.g. 48-80 trees per acre). However, the basics behind the modified central leader pruning style are similar for hedgerow and traditionally spaced orchards. A current trial at Nickels Soil Lab in Arbuckle is looking at three different styles of pruning in a hedgerow Chandler orchard; heavily pruned, minimally pruned, and no pruning. Minimal pruning is what we have recommended for years for most lateral bearing walnut varieties. The heavily pruned style is not recommended for most varieties but is seen in some orchards. The no pruning style (or lack of pruning) is very new. It has been getting a lot of attention from growers and researchers alike. This article summarizes our findings and discusses factors that should be taken into account when choosing a pruning style for your next orchard.

### The Trial Summary

The objective of this trial is to consider different training styles for hedgerow Chandler walnuts with the goal of creating strong trees, early yields, reduced pruning cost, and prolonging the time to the first hedging. The trial was planted in 2008 and is going into its fifth growing season. The trees are planted on berms at a 15 x 22 ft. spacing. The soil is Hillgate loam and Arbuckle sandy loam. The trial is a comparison of three styles of pruning; heavily pruned, minimally pruned and unpruned or no pruning. They are described in detail in the table on the next page. The simplest way to describe the difference between heavily pruned and minimally pruned is that the heavily pruned has branches removed if they are competing with selected scaffolds and all branches are headed in the first

three years. In minimally pruned trees, the competing branches are typically left in the canopy but not headed to develop fruitwood and the chosen scaffolds are headed in years two through four. In both pruning systems, the leader is headed and left as the longest limb.

At this point of the trial the orchard is not yet at full canopy, but light interception data shows no significant difference between treatments except the minimally pruned treatment was slightly larger when measurements were taken in July. The heavily pruned treatment had statistically less crop in the second year of harvest and less cumulative yield for the last two years compared to the minimal and unpruned treatments. Visual observations at this point show more fruiting wood developed in the unpruned and the minimally pruned trees. Unpruned trees produce extension growth every other year with fruiting spurs forming on this extension growth in the following year. Yield efficiency (yield per unit canopy light intercepted) is highest in the unpruned treatment. This means that more nuts are being produced for the size of canopy formed in the unpruned trees. Data from a previous 7 year long trial of pruning versus no pruning on Howard walnut ([http://walnutresearch.ucdavis.edu/2010/2010\\_117.pdf](http://walnutresearch.ucdavis.edu/2010/2010_117.pdf)) showed that although the yield in the early years tended to be higher for unpruned trees, by the time the trees had filled in their allotted space, there were no differences in cumulative yield or quality for the pruned and unpruned treatments.

### **Take Home Lesson**

This trial is not complete nor have we answered some very important questions. We do not know how the pruning treatments will affect limb breakage when the trees come into full bearing. We also do not know if the same response would be observed on different soil types, tree spacing, or irrigation methods. A side trial containing the same treatments on different varieties; Gillet, Forde, and Tulare, is showing similar trends in yield and growth. We also have not answered the question: ‘will an unpruned orchard need to be hedged earlier or later than the other pruning treatments?’ If you are interested in trying a new training/pruning style in your own orchard it would be a good idea to start small and see how your orchard practices work with a more minimal style of pruning. It has been found that once you start a pruning style you should continue with the same style until the trees are mature. Changing midstream is not recommended.

### **Factors that may affect the pruning style you choose**

- Variety of walnut and rootstock: more vigorous scions and rootstocks can create more problems with blind wood and excessive in-season branching with heavy pruning.
- Soil and irrigation: vigor is also affected by soil and water management practices.
- Tree spacing: ideally your overall management (pruning, irrigation and nutrition) should encourage the tree to fill the space allotted without excessive crowding. We may find that higher yield efficiency and more fruiting wood in the early years will be better in the long run. Stay tuned for the answer.

### **Notes on the Forde Variety**

The new Forde variety has been found to be a very vigorous tree and in our pruning trial the heavily pruned Forde trees had many of the side buds break in the current season and form groups of thin branches, also known as “witches broom”. These problems have also been observed by a number of growers. At this point we suggest only minimal pruning for Forde with no heavy pruning or heading of the scaffolds. (Full report at <http://walnutresearch.ucdavis.edu>)

### **Notes on the Chandler Variety**

Although the research trial in Arbuckle is a Chandler hedgerow, Chandler’s vigor, growth and fruiting habit is typically best suited to planting it in a traditionally spaced orchard for long term productivity. What we are learning on growth response to pruning or no pruning during the training stage however, applies to Chandler growing in either configuration.


### Description of the Three Training/Pruning Styles Compared in the Chandler Hedgerow Trial

	Heavily Pruned	Minimally Pruned	No Pruning (or heading)
<b>At planting 2008/first growing season</b>	Trees in all treatments were planted and headed at 4-5 healthy strong buds above graft union on the scion. One strong leader was selected for the trunk and a few lower shoots above the graft union were kept shorter than the leader but allowed to grow longer than is typical to take some vigor from the unstaked leader to avoid breakage.(note: if trees had been traditionally spaced, they would have been staked)		
<b>First year Dormant pruning 2009</b>	At the end of the dormant season in March, the leader was headed at 6 feet. Lower limbs below 3 feet were removed.	At the end of the dormant season in March, the leader was headed at 6 feet. Lower limbs below 3 feet were removed.	At the end of the dormant season in March, the leader was <b>not</b> headed. Lower limbs below 3 feet were removed. A long stake was placed to support the unpruned leader.
<b>Second year delayed dormant pruning 2010</b>	Selected and headed a central leader by 1/3 of length of current growth. Selected 4-6 primary scaffolds and headed each by 1/3. Heights of scaffolds were maintained below the height of the central leader. All unselected branches were removed if they were likely to compete with chosen scaffolds. Forked branches were reduced to a single branch. All remaining branches were tipped or removed. Lowest branch left at about 4 feet above berm in the row.	Central leader selected and 1/3 of current growth was removed. 4-6 primary scaffolds were selected and headed below the height of the central leader. These heading cuts removed 1/4 to 1/3 of the length of current growth. Forked branches on chosen scaffolds were reduced to a single branch. Remaining unselected branches and small caliper fruit wood were left unpruned and unheaded to create early fruiting wood. Lowest branch left at about 4 feet above berm in the row.	No pruning or heading unless lower branches needed to be removed for reasons of safety or ease of maintenance and harvest.
<b>Third year delayed dormant pruning 2011</b>	The central leader from previous year was examined and if choked out by a stronger branch, a new central leader was chosen. The chosen central leader had 1/3 of the growth from current year removed. Branches that were competing with the leader were removed. In season branching points were removed in secondary scaffolds. Secondary scaffolds were chosen in all directions and headed by 1/3 making sure no secondary scaffold was taller than central leader. All twisted, crossed or rubbing branches were removed.	The strongest, tallest scaffold was chosen as the leader and 1/3 of the current growth was removed. Other scaffolds were left alone if they were growing in a vertical position. One or two strong scaffolds were chosen on the sides of the canopy, one in each cardinal direction was headed removing 1/3 of the current growth. Forked branches were left but twisted, rubbing or overlapping branches were removed.	No pruning or heading unless lower branches need to be removed for reasons of safety or ease of maintenance and harvest.

# Tools for Weed Management Decisions in Orchard Crops

*Brad Hanson, UC Extension Weed Specialist, Davis*

Weed control issues and management tools sometimes seem like a constantly moving target. Weed species shift as orchards age, new species sometimes appear, and herbicide resistant species have occurred in some cases. Similarly, herbicide options can vary significantly among crops and occasionally vary year-to-year as new herbicides are registered or old material are no longer supported by the manufacturer or allowed for use in certain crops. Fortunately, a few tools are available to help California crop producers make informed weed control decisions.



**Weed ID:** First, to make an informed weed control decision, it is important to understand what weed species you are trying to control. One easy (and free) online resource for identifying weeds is available at the University of California Weed Research and Information Center (<http://wric.ucdavis.edu/>). From the main page, go to the “quick links” on the right side of the screen and click on the Weed ID Tool. The first step is to identify if you have a broadleaf or a grass-like weed. Once you’ve selected one of those, you are brought to a series of pull-down menus that allow you to narrow down a list of common weeds based on physical characteristics. This Weed ID Tool is designed for those of us who may not be expert botanists so if you don’t have a plant part, or if you don’t understand the botanical term, you can simply skip that characteristic. Once you’ve made your selections, click the “search database” button to look at the narrowed down list of plants remaining in the database.




California Herbicide Registration on Horticultural Tree and Vine Crops - (updated January 2012) - UC Cooperative Extension

Herbicide-Common Name (example trade name)	Site of Action Group <sup>1</sup>	Almond	Pecan	Pistachio	Walnut	Apple	Pear	Apricot	Cherry	Nectarine	Peach	Plum / Prune	Avocado	Citrus	Date	Fig	Grape	Kiwi	Olive	Pomegranate
<b>Prememergence</b>																				
dichlobenil (Casoron)	L / 20	N	N	N	N	R	R	N	R	N	N	N	N	N	N	N	R	N	N	N
diuron (Karmex-Diurex)	C2 / 7	N	R	N	R	R	R	N	N	N	R	N	N	R	N	N	R	N	R	N
EPTC (Eptam)	N / 8	R	N	N	R	N	N	N	N	N	N	N	N	R	N	N	N	N	N	N
flumioxazin (Chateau)	E / 14	R	NB	R	R	R	R	R	R	R	R	R	NB	NB	N	NB	R	N	NB	NB
indaziflam (Alion)	L / 29	R	R	R	R	R	R	R	R	R	R	R	N	R	N	N	R	N	N	N
isoxaben (Trellis)	L / 21	R	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	N	NB	R	NB	NB	NB
napropamide (Devrinol)	K3 / 15	R	N	N	N	N	N	N	N	N	N	N	N	N	N	N	R	R	N	N
norflurazon (Sollicam)	F1 / 12	R	R	R	R	R	R	R	R	R	R	R	R	R	N	N	R	R	N	N
oryzalin (Surflan, Farm Saver)	K1 / 3	R	R	R	R	R	R	R	R	R	R	R	R	R	N	N	R	R	N	N
oxyfluorfen (Goal, GoalTender)	E / 14	R	R	R	R	R	R	R	R	R	R	R	R	NB	N	N	R	R	R	R
pendimethalin (Prowl H <sub>2</sub> O)	K1 / 3	R	R	R	R	R	R	R	R	R	R	R	R	R	N	N	R	R	R	R
penoxsulam (Pindar GT)	B / 2	R	R	R	R	R	R	R	R	R	R	R	R	R	N	N	R	R	R	R
pronamide (Kerb)	K1 / 3	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	R	R	N	N
rimsulfuron (Matrix, Mana)	B / 2	R	R	R	R	R	R	R	R	R	R	R	R	R	N	N	R	R	N	N
simazine (Princep, Caliber 90)	C1 / 5	R	R	R	R	R	R	R	R	R	R	R	R	R	N	N	R	R	N	N
thiazopyr (Visor)	K1 / 3	NB	N	NB	NB	N	N	NB	NB	NB	NB	NB	N	R <sup>2</sup>	N	NB	R	N	R	N
<b>Postemergence</b>																				
carfentrazone (Shark, Rage)	E / 14	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
clethodim (Prism)	A / 1	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	N	R	N	NB	R	N	NB	N
clove oil (Matratec)	NC <sup>3</sup>	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
2,4-D (Clean-crop, Orchard Master)	O / 4	R	R	R	R	R	R	R	R	R	R	R	NB	N	N	NB	R	N	N	N
diquat (Diquat)	D / 22	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	N	NB	R	N	NB	NB
d-limonene (GreenMatch)	NC <sup>3</sup>	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
fluzafop-p-butyl (Fusilade)	A / 1	NB	R	R	R	R	R	R	R	R	R	R	NB	R	N	NB	R	N	NB	N
glyphosate (Roundup)	G / 9	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
glufosinate (Rely 280)	H / 10	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
halosulfuron (Sandea)	B / 2	N	R	R	R	R	R	R	R	R	R	R	N	N	N	N	R	N	N	N
paraquat (Gramoxone Inteon)	D / 22	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
pelargonic acid (Scythe)	NC <sup>3</sup>	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
pyraflufen (Venue)	E / 14	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
safinacil (Treevix)	E / 14	R	N	R	R	R	R	R	R	R	R	R	N	R	N	R	R	R	R	R
sethoxydim (Poast)	A / 1	R	R	R	R	R	R	R	R	R	R	R	NB	R	NB	R	R	R	NB	NB

Notes: N = Not registered, NB = nonbearing, R = Registered. This chart is intended as a general guide only.  
 Always consult a current label before using any herbicide as labels change frequently and often contain special restrictions regarding use of a company's product.  
 Weed susceptibility information and the most up to date version of this table can be found at the Weed Research and Information Center (<http://wric.ucdavis.edu>)  
<sup>1</sup> Herbicide site of action designations are according to the Herbicide Resistance Action Committee (letters) and the Weed Science Society of America (number) systems.  
<sup>2</sup> Simazine is registered on only sour cherry in CA. Thiazopyr is registered on orange and grapefruit only.  
<sup>3</sup> NC = no accepted site of action classification; these contact herbicides are general membrane disruptors.

Also available online at: <http://ucanr.org/sites/wric3/?blogpost=6524&blogasset=32026> (UCD Weed Science blog post 1-15-2012)  
 or [http://wric.ucdavis.edu/PDFs/herbicide\\_registration\\_on\\_horticultural\\_tree\\_and\\_vine\\_%20crops\\_2012\\_01.pdf](http://wric.ucdavis.edu/PDFs/herbicide_registration_on_horticultural_tree_and_vine_%20crops_2012_01.pdf) (UC Weed Research and Information Center)



# Walnut Pruning Field Meeting

March 6th, 2012

9:30 a.m. - 12:00 noon

Location: Nickels Soil Lab, Arbuckle

Demonstration of pruning 4 year old hedgerows and conventional planted walnut with a discussion on the pruning trials for hedgerow Chandler.

We will also look at different pruning treatments on Forde and Gillet

The workshop will be conducted by UC Farm Advisors:

Janine Hasey, Sutter & Yuba Counties; Carolyn DeBuse, Yolo & Solano Counties;  
Bruce Lampinen, Pomology Specialist, UC Davis; John Edstrom, Colusa County

IN THE EVENT OF RAIN, MEETING WILL BE HELD ON MARCH 13<sup>th</sup>, 9:30-12 noon

Questions: Yolo UCCE OFFICE AT 530-666-8143 or look at the websites for updated

Information at [ceyolo.ucdavis.edu](http://ceyolo.ucdavis.edu) or [cesolano.ucdavis.edu](http://cesolano.ucdavis.edu)

**Map inside.**

Glenn County  
Cooperative Extension  
P. O. Box 697  
Orland, CA 95963