

*In this issue:*

- Walnut Blight Management
- Flooded Orchards: What To Do
- Monitoring Codling Moth In and Near Mating Disruption Orchards
- IPM Breakfast Meetings Announcement
- Managing Bot Canker & Blight: Research Updates
- Orchard Management Considerations: Budbreak to Early Summer
- Field Meeting

*Submitted by:*

Dani Lightle,  
UCCE Advisor

## Walnut Blight Management

*Richard P. Buchner, UCCE Farm Advisor Tehama, Glenn and Butte Counties*

Walnut blight is the result of infection by the bacterium *Xanthomonas arboricola* pv *juglandis* (*Xaj*). All green tissue is susceptible to infection, including buds, flowers and leaves but nut infections are the most serious, usually resulting in economic damage. Early in the spring, infected walnuts develop a dark sunken lesion at the blossom end ("end blight") killing the developing kernel. As walnuts mature, blight lesions can develop elsewhere on the husk surface ("side blight"). Husk lesions begin as small water-soaked spots that later darken, enlarge, sink and often crack. Infections that do not invade the kernel may increase the possibility of secondary insect or disease damage. Drought years do not favor infection but years with extensive spring rainfall can be disastrous if management action is not taken.

The walnut blight pathogen, *Xaj*, survives the winter in the outer bud scales of dormant buds. As the new shoot elongates after bud break, *Xaj* are water transported from the outer bud scales to green tissue, flowers and developing nuts. Disease control requires applying spray materials to protect green tissue. A successful walnut blight control program focuses on protecting developing shoots and flowers and decreasing *Xaj* bacteria over-wintering in dormant buds.

**Disease Control.** The probability of infection depends upon how much pathogen exists on individual buds and environmental conditions favoring bacterial spread and infection. First blight sprays are timed to coincide with early shoot emergence. This places a protective layer of bactericide on emerging green tissue. Since all shoots do not emerge at the same time, a good compromise is to apply the first spray application when 40% of the shoots on the tree are at the "prayer stage" when unfolded leaves resemble hands held in prayer. A second spray is applied about 7 to 10 days later to protect the remaining opening buds. Additional sprays may be necessary depending upon the initial inoculum, disease history, weather conditions and variety. It is possible that half spray programs favor increasing populations of *Xaj* and the half spray-sub lethal exposure ap-

**ANR NONDISCRIMINATION AND AFFIRMATIVE ACTION POLICY STATEMENT:** It is the policy of the University of California (UC) and the UC Division of Agriculture and Natural Resources not to engage in discrimination against or harassment of any person in any of its programs or activities (Complete nondiscrimination policy statement can be found at <http://ucanr.edu/sites/anrstaff/files/169224.pdf>). Inquiries regarding ANR's nondiscrimination policies may be directed to Linda Marie Manton, Affirmative Action Contact, University of California, Agriculture and Natural Resources, 2801 Second Street, Davis, CA 95618, (530) 750-1318.

To simplify information, trade names of products may be used. No endorsement of named products is intended, nor is criticism implied of similar products which are not mentioned.





40% of shoots at “prayer stage” is the appropriate timing for the first walnut blight spray.

proach is an excellent way to select for resistance to the only effective spray program we have for walnut blight.

**Material Choice.** Copper formulations tank mixed with mancozeb are the best currently available spray materials for walnut blight management. Resistance to copper alone is very common in California and is thought to be responsible for

poor walnut blight control in the early 1990's prior to EBDC tank mixes. Under heavy blight pressure, using maximum rates allowed by the label is suggested. In February, 2017 the Department of Pesticide Regulation approved requests by multiple County Ag Commissioners for emergency application of fungicides to orchards with standing water. The end date is June 1, 2017. Contact your County Agricultural Commissioner's office in Butte, Colusa, Glenn, Merced, San Joaquin, Stanislaus, Sutter, Tehama, Yolo and Yuba Counties for additional information.

### Some Reasons Walnut Blight Management Fails

1. First spray timing too late.
2. Walnut blight bacterial population increases in dormant buds resulting in high initial disease pressure.
3. Material rates too low.
4. Poor spray coverage by air or ground.
5. Using a weak bactericide in high blight potential orchards.
6. “Half sprays” due to every-other-row application.
7. Not tank mixing copper compounds with a Mancozeb formulation.
8. Dense tree canopies.

## Flooded Orchards: What to Do in 2017

*Janine Hasey, UCCE Farm Advisor, Sutter/Yuba/Colusa Cos.  
Greg Browne, USDA Plant Pathologist, UC Davis*

Experiences with the 1986 and 1997 levee breaks in Yuba and Sutter Counties and other “high-water” events over the past thirty years afford us some expectations for the types and extents of tree damage that may result from orchard flooding in 2017. This article will first discuss some generalities of flooding damage and two key types of damage that tend to follow prolonged orchard flooding, i.e., waterlogging and *Phytophthora* diseases. Next, the article reflects on the floods in 1986 and 1997 and considers ways in which specific conditions associated with those floods seemed to affect orchard outcomes. Finally, we offer management steps to consider in dealing with orchard flooding in 2017.

**Flood Damage Generalities.** Tree damage from flooded soil is usually minimal if the flood occurs when the trees are dormant and the water continues to flow rather than remains standing in the orchard area. However, where water stands for extended periods in an orchard, two potential problems are of concern: 1) waterlogging of the root system, and 2) diseases caused by *Phytophthora*.

**Waterlogging.** Waterlogging can be viewed as an “oxygen starvation” effect of flooding. When an orchard soil is flooded or otherwise saturated with water, the air in its pore spaces is displaced, removing the supply of oxygen to the roots. The terminal portions of new walnut roots can be killed within one to four days in a saturated soil. As saturation is prolonged, damage to roots becomes greater and can lead to chlorosis, leaf wilting, and in extreme cases, entire root systems can be killed. As soil temperatures increase, the negative effects of waterlogging are accelerated. Walnut trees apparently survive waterlogging events during winter due to a combination of low soil temperature and a lack of active roots.

Waterlogging risk tends to be greater in “heavy” or compacted soils because of slower reentry of oxygen after saturation. Following floods in previous years, we observed less injury to trees growing in deep, permeable soils than to trees growing on soils with impaired vertical drainage caused by a claypan or hardpan.

Waterlogging damage to walnut trees can vary in severity, may not be immediately manifest, and can be difficult to discriminate from *Phytophthora* damage. For example, decline of walnut trees is sometimes not obvious until late spring or summer, when backhoe excavations can be used to confirm suspected waterlogging damage on walnut root systems.



Swollen lenticels (white raised areas) on clonal Paradox

Boundaries between dead, waterlogged portions of major roots and crowns and living parts of the same roots tend to be poorly defined and occur in the absence of distinct cankers and lab diagnostics that indicate *Phytophthora* crown and root rot. Waterlogged tissues often have swollen lenticels.

**Phytophthora diseases.** Several species of the “water mold” *Phytophthora* are found in surface water sources such as our rivers and canals, as well as in infested soils. These fungal-like organisms can be carried along by the water and swim in it. After a levee break, the river water is a potential source of spreading *Phytophthora* inoculum throughout the



Aerial *Phytophthora* bleeding cankers (arrow) and tissue damage (evident where bark removed). Photo: J. Hasey

flooded area. Flooding can bring active *Phytophthora* inoculum in contact with the aerial portions of trees, resulting in infections of the tree trunk, scaffolds and shoots. Flooding also increases the risk of root and crown infections by *Phytophthora*. Some *Phytophthora* species preferentially infect roots, but many preferentially infect the root, crown, or aerial tree parts.

When *Phytophthora* infects woody tissues of trees above or belowground, it typically causes cankers (i.e., continuous, often-elliptical areas of dead bark, ranging from less than an inch to more than a foot across). The cankers often “bleed”. Removing the outer bark from the margin of a canker typically reveals a distinct, zonate (with concentric lines) margin, which is in contrast to the typically diffuse, non-zonate margin and vascular streaking evident at the edges of dead areas of waterlogged roots.

In general, cool to moderate temperatures and water-saturated conditions in or on soil favor tree infection by *Phytophthora* species. Some *Phytophthora* species are more virulent than others, and risk of disease also depends on genetic resistance of the rootstock or scion. Within a given rootstock or scion, susceptibility to *Phytophthora* can vary with time of year. For example, in experiments on seasonal variation in susceptibility of walnut, almond, and peach rootstocks to *Phytophthora citricola* (which can cause root and crown rot and trunk and branch cankers), the root-

stocks were least susceptible to infection during wintertime and most susceptible during the growing season. In contrast, infection by *Phytophthora syringae* in almond and peach is most likely during the cool conditions of fall, winter, and early spring. These cankers cease development when summer temperatures become lethal to this "cool-temperature" pathogen.

### 2017 flooding and management considerations

**Flood overview.** From a series of tropical storms, we have experienced record-breaking rainfall events in January and February which have resulted in very high river flows for many weeks. At this writing, orchards **outside the levees** along the rivers either are or have been flooded from river seepage due to high flows and/or overflowing ditches/canals. With the record-breaking snow pack, there is uncertainty as to how long the rivers will have high flows or if we have a flood event during springtime. It likely will take months before the extent of losses from waterlogging and/or disease can be assessed. As the trees become active and the weather warms up, we can expect to start seeing losses that could continue through the hot summer when trees with damaged root systems or crowns typically collapse. From recent research, substantial root growth in walnuts begins about a month after leaf-out and peaks in the summer. This should work in our favor for orchards that are flooded or saturated this spring.

### Management guidelines to consider:

- Drain or pump standing water out of orchards.
- Reduce vehicle/farm equipment traffic. Wet soils are easily compacted. Delay all operations that can wait until soil is dry enough to crumble at a depth of five to six inches rather than slick over or pack.
- Apply sprays by air. The Department of Pesticide Regulation issued the emergency application of

several fungicides to orchards with **standing water only** including Butte, Colusa, Glenn, Sutter, Tehama, Yolo, and Yuba Counties This will be in effect until June 1, 2017. See your local Ag Commissioner for more information on allowable fungicides.

- Apply a phosphonate spray in May where *Phytophthora* is found or suspected. A summer and early fall application may also be needed.
- Ridomil application is another option but is considered by some to be less cost effective than phosphonates for some *Phytophthora* diseases.
- Remove deposited silt and debris around the root crowns when possible to decrease the chances of root/crown decay.
- Plants are effective in drying waterlogged soils. Encourage the growth of cover crops or even weeds that will help dry the soil after flooding.
- Fill in eroded areas in orchards if soil is available. Deposited materials in many instances are beneficial.
- Check for salts (chloride and sodium); a continued high water table saturating surface soil may result in these salts accumulating or alkalization of certain spots which have shown these troubles before. Neither leaching nor gypsum treatments will be effective until the water table is lowered and good drainage can be achieved.



Flooded walnuts in March 1986. Photo: J. Hasey

- For new orchards or replants where seepage is problematic, consider using clonal Paradox RX1; it has high resistance to *Phytophthora*, but more observations are needed to determine how it performs under prolonged waterlogging.

**Additional resources.** For additional resources please see Hasey's blog posts at: <http://www.sacvalleyorchards.com/blog/walnuts-blog/> The blog covers several flooding-related issues not covered here, including reflections on previous flood events, guidance on documenting and reporting damage and losses, and locating possible disaster relief resources for flooded orchards outside and inside the levees, including riverbank sloughing, such as occurred extensively along the Feather River.

### **Monitoring Codling Moth In and Near Mating Disruption Orchards**

*Emily J. Symmes, Sacramento Valley Area IPM Advisor  
UCCE Statewide IPM Program*

It is important to understand how mating disruption affects the ability to monitor codling moth in walnut orchards under mating disruption (MD) as well as in orchards in proximity to MD. Researchers are currently investigating just how far-reaching the impacts of mating disruption are to nearby non-MD blocks. Pheromone-only trapping methods will certainly be impacted in MD blocks, and can be affected in nearby non-MD environments (without the added benefit of substantial disruption, and thus population and damage reduction). Other monitoring methods are necessary in and near MD orchards to effectively manage codling moth populations.

Disruption of sexual communication in moths is thought to function by the following broad types of behavioral mechanisms, and combinations of the two mechanisms:

1) Competitive attraction (false-plume-following), in which males are diverted from orienting to females because they are attracted to competing 'false' trails

emitted by synthetic pheromone dispensers.

2) Non-competitive mechanisms, whereby exposure to synthetic pheromone inhibits or blocks the ability of males to sense and/or respond normally to pheromone emitted from females. These include camouflage, desensitization (i.e., adaptation and habituation), and sensory imbalance.

In a practical sense, the presence of synthetic pheromone in and around the orchard environment, if effectively impacting mate location/mating success, will impact our ability to use pheromone lures to track population abundance and activity. Codling moth (CM) females emit sex pheromones which elicit responses from males of the species. Therefore, pheromone traps predominantly track only male activity. Successful MD will shut down pheromone traps to zero or very low male catches relative to non-MD environments. In a MD orchard, you will want to monitor pheromone-only traps to gauge this measure of MD effectiveness. However, it is also critical to be able to track flights and relative population abundance to determine if supplemental insecticide treatments are necessary, and to properly time the applications.

Fortunately, there are options available for monitoring CM in/near MD that can overcome trap shut-down by (1) incorporating non-pheromone based lures to trap males, females, or both sexes; and (2) employing additional surveillance methods (non-trap-based) to gauge pest pressure and inform treatment decisions.

Combination lures:

- Both female and male moths respond to pear ester. This is available in the CM-DA combo lure (contains codlemone, the codling moth pheromone, plus pear ester, a plant volatile-based kairomone). Traps baited with these lures should be used in orchards under MD in addition to pheromone only traps (baited with 1X or L2 lures), necessary to detect trap shutdown as a proxy for

efficacy of the MD treatment. If you are concerned with the performance of pheromone-only CM traps in non-MD orchards, consider adding some CM-DA traps, particularly if you suspect you are in proximity to an MD block that is affecting your pheromone trap catches.

- Another option is the three-way lure (CM-DA combo plus acetic acid, AA). Think of this as a ‘super-charged’ lure, which can provide more robust capture in terms of numbers of males and females (in both MD and non-MD orchards).
- With both the CM-DA and CM-DA+AA lures, be cautious in your evaluation of trap numbers. Numbers may be higher or lower than pheromone trap catch numbers, depending on your particular environment (MD, non-MD, near MD), and may not directly correlate with the same moths/trap/night thresholds you are accustomed to.

Non-trapping methods:

- These surveillance methods involve in-season damage/population estimates via dropped nut counts and canopy counts looking for strikes (visible as small chimneys of frass at the CM entry point). These monitoring methods can and should be incorporated into your IPM program to aid in treatment decisions, regardless of presence of MD and the trap-lure combination you are using.

More information and details regarding these monitoring and treatment options for codling moth are available online in the UC IPM Guidelines ([ipm.ucanr.edu/PMG/r881300211.html](http://ipm.ucanr.edu/PMG/r881300211.html)).

Best practices for lure-based trapping and monitoring in general:

**Storage and handling.** Always follow manufacturer guidelines for storage and handling of lures. Most

lures are best kept refrigerated or frozen (in non-cycling freezers). It is best to order fresh lures each year, as keeping lures for multiple years can impact performance. If lures are particularly ‘hot’ initially, pre-age lures (open packaging and let them sit out for a day or so) prior to deploying in the field to avoid misinterpretation of activity peaks. Use new traps every year and wear disposable rubber gloves when handling lures and traps to avoid cross-contamination (this will help prevent by-catch of non-target species).

**Interpreting catches and perceived failures.** If you are encountering trap catch data that seems unusual, first consider any potential storage, handling, or contamination issues. Next consider other factors, such

### Upcoming IPM Breakfast Meetings

Join Area IPM and Farm Advisors to discuss current pest management and production issues. We will largely focus on orchard crops (but everything is on the table for discussion!). These meetings are open to all interested growers, consultants, PCAs, CCAs, and related industry.

Meetings will be held the second Tuesday of each month through November and will cover a wide range of timely pest and orchard management topics. Meeting locations will be rotated throughout the Sacramento Valley each month. Please contact Emily Symmes ([ejsymmes@ucanr.edu](mailto:ejsymmes@ucanr.edu)) to request topics or bring your questions to the meeting!

#### Upcoming meetings:

- Glenn: May 9<sup>th</sup>
- Butte: June 13<sup>th</sup>
- Colusa: July 11<sup>th</sup>

Remaining 2017 meetings will be held in August (Yuba-Sutter), September (Tehama), October (Glenn), and November (Butte).

Additional information for each meeting will be available on the events page at:

## Managing Bot Canker and Blight in Walnut - 2016 Research Updates

Janine Hasey, UCCE Farm Advisor, Sutter/Yuba/Colusa Cos.

Themis Michailides, Plant Pathologist, UC Davis/Kearney Research and Extension Center, Parlier

Rainfall favors *Botryosphaeria* (Bot) and *Phomopsis* canker and blight of walnut when temperatures are over 50° F, so be alert to disease spread this spring and summer. Bot reduces yields by killing small fruit wood and large branches and directly infecting the nut. It is most obviously seen in walnut orchards as blighted shoots (dead branches with the leaves still stuck on), blighted twigs (dead, darkened and shriveled) and fruit with the entire hull blackened but still on the tree. Fruit can also have early season latent infections, but tissue death won't be visible until late summer/early fall. Both pathogens (*Botryosphaeria* & *Phomopsis*) produce abundant water-splashed pycnidiospores and wind-borne ascospores. Practically-speaking, this means Bot can spread by spores dripping and splashing OR by blowing around in the orchard. Both spore types need water to trigger infection.



Spurs killed by Bot or Phomopsis. These will be evident during leaf-out. Photo: T. Michailides.

**Conditions needed for disease.** Bot infection requires susceptible tissue, presence of infective Bot spores, and the right environmental conditions - at least ¼" of rain (or irrigation water directly on susceptible tissue) and temperatures at or above 50°F.

**Disease management practices** integrate cultural (pruning) and chemical control.

- Cultural control: reduce inoculum by pruning out dead branches, branch stubs, and blighted shoots in the summer or fall when not raining; disinfect pruning equipment; avoid sprinkler irrigation that wets the canopy.
- Chemical control: apply effective fungicides at proper timing adjusted for weather and Bot inoculum level in orchard. There are several fungicides that have reduced blighted spurs compared to the control in studies and their efficacy can be seen at <http://www.ipm.ucanr.edu/PDF/PMG/fungicideefficacytiming.pdf> in the walnut section.

### What's New - 2016 Research Update

**Fungicide spray timing** based on rain and temperature delivered the best control. This may be the most important finding if we continue to have a wet spring: Using the Leaf Wetness Model (LWM), sprays were only applied when there were environmental conditions conducive to disease ( $\geq 1/4$ " rain and temperatures  $\geq 50^\circ$  F). This resulted in three sprays in both 2015 and 2016. A fungicide is applied within 2-3 days of a rain event and estimate a 2 week residual.

- Use the LWM graph in Figure 1 (next page) during a rain event to pinpoint the BOT risk zone and deter-

**Relative risk of infection of pistachio or walnut by *Botryosphaeria* as affected by wetness period (W) events and Temperature (T)**





Figure 1. Based on rainfall amount and temperature, use this LWM graph during a rain event to pinpoint the BOT risk zone and determine whether a spray is needed. A spray is applied when points fall in medium and high risk. If a point falls on the line separating low and medium risk, a spray is also applied

mine whether a spray is needed.

**2016: Leaf Wetness Model (LWM) where rain exceeded threshold in Chandler (Butte Co.)**

- Figure 2 (right) shows how rain events triggered three sprays in 2016. The only reason the high rainfall, high risk event on May 5-7 was not sprayed is because it was within the 2-week fungicide residual period of an earlier spray. Note that if a point falls on the line separating low and medium risk, i.e. during the rain event on May 21, a spray is also applied.



Sprays: 26 April, 23 May, 20 June 2016

Figure 2. In 2016, the LWM triggered four risk events but only three sprays were needed. The high risk event fell in the period when there was residual fungicide activity.

- The LWM was more economical than the 4 calendar sprays that gave equal control:  
 April (bloom)+May+June+July  
 or  
 May+June+July+November (post-harvest).

**If only applying one fungicide spray**, the second part of June to first part of July had the most effect, when tested in both a low (8% infection when unsprayed, 2015) and high (38% infection when unsprayed, 2016) disease year in the same orchard.

- When sprayed just once at either bloom, mid-May, mid-June, mid-July, mid-August or post-harvest (October or early November), both the **mid-June** and **mid-July sprays** significantly decreased blighted shoots when compared with no spray treatment.

**Bot inoculum and brush disposal/spray scenarios:** Chipping pruned wood reduces ability to cause infection by only two-thirds, so brush removal is still best option in light to moderate infected orchards. Always prune or hedge your more lightly infected orchards before moving to heavily infected ones. Watch for rains and spray according to leaf wetness model if possible.

- *Heavily infected orchards (>50%):* Chip prunings and may leave in orchard since already high inoculum. Use a full spray fungicide program. (From experience, when inoculum on the trees is very high, there is a saturation point beyond which additional spore loads reaching the canopy do not cause more infection).
- *Moderate (20-50%) infected orchards:* Do not chip prunings but remove from the orchard and burn them if permitted. Some fungicide spraying.
- *Light (1-20%) infected orchards:* Do not chip prunings but remove from the orchard and burn them if permitted. Use one mid-June or early to mid-July spray.
- *No Bot infection (mostly only applies to young orchards):* Chip prunings and leave in orchard. No sprays are needed.

## Orchard Management Considerations: Budbreak through Early Summer

*Katherine Pope, UCCE Orchard Advisor, Sacramento, Solano and Yolo Counties*

### APRIL

- Stay on top of blight sprays, especially if we have rain around “prayer stage”. See article (this newsletter).
- The first flight of codling moth typically starts between mid-March and mid-April and will have two peaks (1A and 1B). Male biofix is the first day male moths are consistently found in traps and sunset temperatures reach 62°F. In orchards not using mating disruption with more than 3% damage last season, treating 1A and 1B flights may be warranted if pheromone traps (1X or L2) are catching more than an average of 2 moths/trap/night. Treat 1A larvae about 250 to 300 DD after biofix, depending on material. Some newer “reduced-risk” chemistries are more effective applied on the early end. If 1A treatment is not needed based on orchard history and/or trap catches, delaying treatment until 1B or second flight will minimize interference with walnut aphid parasitoids. For more information, see <http://ipm.ucanr.edu/PMG/r881300211.html>.
- Consider putting out navel orangeworm (NOW) pheromone traps (Biolure or L2), at least in more susceptible varieties and/or blocks with a history of NOW damage. See Winter 2016/17 NOW Update article for more information: [http://cesutter.ucanr.edu/newsletters/Sacramento\\_Valley\\_Walnut\\_News66483.pdf](http://cesutter.ucanr.edu/newsletters/Sacramento_Valley_Walnut_News66483.pdf)
- Limbs that have been killed by Bot canker are easy to identify between budbreak and full leaf expansion. But wait to prune out infected limbs once rain is no longer forecasted, i.e. early summer. For more on Bot, see the article in this newsletter.

- If timing Bot treatment based on the Leaf Wetness Model, watch for storms that bring  $\geq \frac{1}{4}$ " rain and temperatures  $\geq 50$  F. See the Bot article in this newsletter for more on treatment timing.
- For orchards with standing water, take pictures to document the flooding. This will help with diagnosing problems later in the season and is needed if applying for disaster relief programs. Photos taken on your phone will usually have a "created" file date that will help with record keeping. For more, see the article on flooded orchard damage in this newsletter.
- Apply foliar zinc (if needed, based on leaf sample analysis) when shoots are 6 to 10 inches long, when zinc can be easily absorbed through the leaf surface. If the deficiency is severe, additional sprays can be applied two more times every 2 to 3 weeks.
- Perform irrigation system maintenance now, before irrigation is necessary and system problems could cause tree stress. Check for broken or clogged filters and emitters. See <http://micromaintain.ucanr.edu/> for more tips on maintaining micro-irrigation systems.
- Scout for scale crawlers by putting out double-sided sticky tape in late April, if scale has been a problem and you didn't treat for scale during the dormant season.

## MAY

- Watch traps for the codling moth 1B peak about 600 to 700 DD after the first biofix. If a significant 1B flight peak occurs (and damage exceeded 3% last season), consider treating when residual effectiveness of the first treatment has elapsed. If a 1A spray was not applied, 1B treatment should be approximately 650 to 700 DD from first biofix (confirm with trap activity; ideal timing depends on material). More at <http://ipm.ucanr.edu/PMG/r881300211.html>.
- Apply the first round of nitrogen fertilizer in May, not before. Walnut trees only use stored nitrogen the first month after leaf-out, meaning N applied before May will likely be leached by rain and/or irrigation. Walnut tree nitrogen use is fairly steady over the growing season. Evenly dividing nitrogen application in 3 -4 doses between May and the end of August will improve N uptake compared to 1-2 applications.
- Apply a fungicide spray for Bot canker and blight around mid-month in orchards with moderate to heavy infection.
- Survey weeds to see which weeds were not controlled by fall or winter treatment. The UC Weed ID Tool at <http://weedid.wisc.edu/ca/weedid.php> can help with identification.
- Aphid sampling should begin this month and continue throughout spring and summer. Collect 5 first sub-terminal leaflets (one back from the last leaflet) from 10 trees, checking the top surface for dusky-veined aphids and the underside for walnut aphids. Make treatment decisions following guidelines at <http://ipm.ucanr.edu/PMG/r881300511.html>.

## JUNE

- Hang Walnut Husk Fly traps by June 1. Yellow sticky traps charged with an ammonium carbonate lure work best. Check traps 2 to 3 times per week and treat based on detection of eggs in trapped females or overall trap catch numbers. For more details on treatment decision-making, see <http://www.sacvalleyorchards.com/walnuts/insects-mites-walnuts/walnut-husk-fly-biology-monitoring-and-spray>



### **Nickels Field Day (May 10) includes Three Walnut Talks**

The annual Nickels Field Day will be Wednesday, May 10 beginning at 8:30. The location is Green Bay Rd, Arbuckle.

Walnut Talk topics are in the early morning session (9-10:30) and include:

- Walnut pruning/hedging
- Bot management
- Thousand cankers research update

There will also be an almond topic in the morning session (irrigation) and four almond-related topics in the late morning session (10:30-noon) including canker diseases, variety selection, and disease management.

A no-host lunch (tri-tip), benefiting the Pierce High School FFA program will follow the talks. More information/detailed agenda will be available at <http://www.sacvalleyorchards.com/>.

PCA/PA and CCA credit hours will be available.

### **Airblast Sprayer Calibration Field Day Set for Tuesday, April 25**

A sprayer calibration workshop, co-sponsored by BASF and Univ of California Cooperative Extension will be held Tuesday, April 25 from 7:30 AM – 1:00 PM at the Nickels Soil Lab (Green Bay site) in Arbuckle.

Light breakfast and coffee at 7:30, program starts at 8 and will end at noon with lunch provided. From 1-3 PM, the instructor (Matt Strmiska from ADAPTIV ([adaptiv.us](http://adaptiv.us))) will be available for questions and demo requests.

Four hours of PCA/PA and CCA hours have been requested.

The morning program will include:

- Overview of equipment and best spraying practices

Nonprofit Organization  
U. S. Postage Paid  
Orland, CA 95963  
Permit No. 63

Glenn County  
Cooperative Extension  
P. O. Box 697  
Orland, CA 95963